

**Lost River Sucker**  
*(Deltistes luxatus)*

**5-Year Review:**  
**Summary and Evaluation**

**U.S. Fish and Wildlife Service**  
**Klamath Falls Fish and Wildlife Office**  
**Klamath Falls, Oregon**  
**2024**

## **5-YEAR REVIEW**

### **Lost River Sucker/*Deltistes luxatus***

#### **GENERAL INFORMATION**

**Species:** Lost River Sucker (*Deltistes luxatus*)

**Date listed:** July 18, 1988

**FR citation(s):** 53 FR 27130

**Classification:** Endangered

**Critical habitat:** 77 FR 73740 – December 11, 2012

#### **Methodology used to complete the review:**

In accordance with section 4(c)(2) of the Endangered Species Act of 1973, as amended (Act), the purpose of a 5-year review is to assess the status of threatened or endangered species to determine if it has changed, and whether the species is classified appropriately. A team of U.S. Fish and Wildlife Service (Service) employees from the Klamath Falls Fish and Wildlife Office (KFFWO) evaluated the status of the Lost River sucker to inform this 5-year review. This team included subject matter experts from KFFWO, the Klamath Falls National Fish Hatchery (KFNFH), and regional staff. The team reviewed the 2019 Species Status Assessment (SSA), the 2019 5-year review, and incorporated the newest scientific information into this review to conclude our determination of the current status of the species. This review represents our evaluation of the best available scientific information, including resource needs and current and future condition of the species.

**FR Notice citation announcing the species is under active review:** 88 FR 56042

#### **REVIEW ANALYSIS**

**Recovery Plan:** U.S. Fish and Wildlife Service. 2012. Revised recovery plan for the Lost River sucker and shortnose sucker (*Deltistes luxatus* and *Chasmistes brevirostris*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California.

**Recovery Criteria:** The recovery criteria comprise a combination of measures that must occur to ameliorate or eliminate threats to the species for each recovery unit to achieve numerical demographic targets. These are found in detail in the Revised Recovery Plan (USFWS 2012). In order to downlist Lost River sucker to threatened status, the following must be achieved:

- A.1 Current spawning and rearing habitat is maintained, and improved access ensures annual use.
- A.2 A range-wide Spawning and Rearing Enhancement Plan has been developed and implemented. This plan shall identify and prioritize areas of potential spawning and rearing habitat for enhancement and/or restoration, including areas which are degraded or

unavailable due to lack of connectivity or passage.

- A.3 Connectivity and access is assured to habitats that provide refuge to suckers to avoid poor water quality (particularly Pelican Bay) during the months of July, August, and September – Upper Klamath Lake (UKL) Recovery Unit. Natural vegetated wetland areas are restored, including in-stream, wetland, and riparian areas around the mouth of Willow Creek where it meets Clear Lake Reservoir and throughout its drainage – Clear Lake Reservoir Management Unit.
- C.1 Newly identified or clarified effects of predation and disease are minimized through implementation of recommendations from ongoing scientific research which clarifies the interaction of Lost River sucker and shortnose sucker with predators and pathogens.
- E.1 An Entrainment Reduction Plan has been developed and implemented. This plan shall identify and prioritize screening of diversions throughout upper Klamath Basin, including the Klamath Project, and propose strategies for efficient reduction of entrainment.
- E.2 Establishment of two additional recurring and successful spring-spawning populations in the Upper Klamath Lake-Spring Management Unit.
- E.3 Development and implementation of a plan to assess, monitor, and improve juvenile and sub-adult vital rates and demography, including threats and negative impact reduction. This plan shall also designate specific demographic or vital rate targets, and strategies for achieving these targets, important for downlisting and delisting.
- E.4 The effects of detrimental water quality have been minimized through implementation of recommendations from ongoing scientific research which clarifies the relationship of these factors with sucker mortality – Upper Klamath Lake Recovery Unit.

In order to delist Lost River sucker the following additional criteria must be met:

- B.1 The States of Oregon and California and the Klamath Tribes, collaboratively or separately, should prepare and finalize population management plan(s) for the Lost River sucker.
- E.5 After 25 years, the average annual rate of population change is greater than one and the number of spawning individuals is greater than what was present in the baseline years for the Upper Klamath Lake River and Upper Klamath Lake Spring Management Units. See Appendix II of the Revised Recovery Plan for descriptions and estimation procedures of these measures. Twenty-five years equates to approximately two average adult life spans for Lost River sucker. This period will enable assessment of the species' response to cyclical threats, such as periodic die-offs and drought. The baseline year for Lost River sucker is 2002, since this is the first year in which estimates of this type are statistically valid for the species.

Of these criteria, only A.3 has been realized through the Interim Plan for Klamath Project Operations 2023 Biological Opinion (USFWS 2023). The Biological Opinion provides guidance to the Bureau of Reclamation that ensures access to areas in Upper Klamath Lake, such as Pelican Bay, under normal operating conditions. Nevertheless, this requires regular reevaluation as conditions and operations change. However, several of these criteria are in process, and are described below; for example criteria C.1, regarding effects of predation and disease on suckers, continues to be investigated by partner organizations through analysis of PIT tags detected on

avian colonies and research into eye fluke parasitism in UKL (Evans et al. 2022, J. Lovy personal comm. 2024).

## **UPDATED INFORMATION**

The brief summary below incorporates updates since the last 5-year review in 2019 that accompanied the SSA (USFWS 2019). Reports on juvenile sucker survival in UKL and Clear Lake remain consistent with little to no recruitment in UKL and high annual variability in Clear Lake (Bart et al. 2020, Martin et al. 2022.). Updated reports were prepared on non-lethal aging techniques, survival, growth, health, and condition of listed juvenile suckers in Upper Klamath Lake and Clear Lake Reservoir, with no changes realized in the past five-years (Hereford et al. 2019, Bart et al. 2020, Burdick et al. 2020a, Burdick et al. 2020b, Markle et al. 2020, Burdick et al. 2021, Hewitt et al. 2021, Martin et al. 2022, Martin et al. 2023).

Multiple mesocosm studies were conducted using juvenile suckers to understand mortality in UKL and to inform net pen rearing operations (Hereford et al. 2019, Burdick et al. 2020, Burdick et al. 2021). One study indicated that greater size at release and location of release can positively influence survival rates of Sucker Assisted Rearing Program (SARP) juvenile suckers in UKL (Caldwell et al. 2023). Additionally, investigations into the effects of disease, parasitism, and predation on survival of wild and hatchery-reared juveniles continues (Markle et al. 2020, Martin et al. 2021a., Evans et al. 2022, J. Lovy personal comm 2024).

Reports on remaining populations reveal a continued decline in the number of spawning adults in the Williamson River and at the shoreline springs in UKL, low recapture probabilities for adult LRS in Clear Lake, and high variability in survival across age-classes in the Lost River Unit (Hewitt et al. 2021, Martin et al. 2021b., Krause et al. 2023). In one study from the UKL Unit, over 600 adult suckers were translocated from Lake Ewauna to UKL after being tagged with a radio telemetry tags to track movement (Banet and Hewitt 2019). The results from this project were limited to only a few years of tracking but indicated that encounters of translocated individuals were two times less likely than resident adult suckers (Banet and Hewitt 2019).

A few important changes were noted for adult suckers in the Lost River Unit, including the lack of LRS encountered during surveys, increasing abundance of non-native species bycatch, and indications that all SNS-like fish captured were KLS or KLS/SNS hybrids based on genetic identification (Martin et al. 2021). An additional update on the Lost River Unit included information on Clear Lake spawning and movement patterns determined through radio telemetry. The reports on PIT tag re-encounter data suggested that LRS remain in the lower portions of tributaries during spawning migrations, while SNS move further upstream (Banet et al. 2021).

Due to the inception of SARP in 2015 and the continued progress with captively rearing suckers, the initial methods for the program were reported in 2021 (Day et al. 2021). A genetic tool to assist staff in sucker identification has been developed using restriction-site-associated DNA sequencing (Smith et al. 2020). This technique provides the capacity to distinguish between the four Klamath Basin sucker species (Lost River sucker, shortnose sucker, Klamath largescale sucker, or smallscale sucker), but only in the UKL Unit (Smith et al. 2020, Smith et al. 2023).

The genetics of LRS indicates that it hybridizes less frequently with other sucker species than SNS (Dowling et al. 2016, Smith et al. 2020, Smith et al. 2023).

Efforts to understand the effects of water quality parameters on LRS and SNS in UKL have also continued. Within the last three years a publication regarding water level fluctuations in UKL and the probability of these fluctuations effecting water quality thresholds for suckers was investigated (Kann and Walker 2020). This study analyzed a long-term data set to evaluate potential relationships between water level fluctuations and pH, un-ionized ammonia concentrations, dissolved oxygen, and chlorophyll-a . The study indicates the probability of exceeding dissolved oxygen, un-ionized ammonia, and pH stress thresholds for suckers increased during both high and low lake elevations . Kann and Walker (2020) therefore recommend maintaining intermediate (but specific) lake levels to decrease the probability of exceeding these stress thresholds for LRS and SNS.

A review was also conducted to collate pertinent information on harmful cyanobacterial blooms and associated effects on juvenile LRS and SNS (Burdick et al. 2020a, and see Burdick et al. 2020b) n this review the effects of cyanotoxins, high pH, hypercapnia, hypoxia, and high levels of ammonia on juvenile LRS and SNS were compiled, unless information was not available at which point other fish species' responses were included (Burdick et al. 2020a). The pH in UKL regularly reached sub-lethal levels for extended periods of time and was noted as a factor that could increase mortality of juvenile suckers (Burdick et al. 2020a). However, there is indication from this review that detrimental water-quality following harmful cyanobacterial blooms and crashes, along with the many other stressors in UKL, can likely cause chronic stress, which can lead to indirect mortality of juvenile suckers (Burdick et al. 2020a). It is important to add that while harmful cyanobacterial blooms persist in UKL, these are largely driven by total phosphorus inflows to the lake during the winter months, and these inflows have been decreasing based on recent analysis of the period 1992-2018 (Walker and Kann 2022).

## **CURRENT SPECIES STATUS**

The following discussion is a brief summary of the interacting influences of physical, chemical, and biological factors that continue to threaten Lost River sucker. To determine whether to list, delist, or reclassify (change from endangered to threatened status, or vice versa) a species under section 4(a) of the ESA, we evaluate five major categories of threats to the species: (A) the present or threatened destruction, modification, or curtailment of its habitat or range; (B) overutilization for commercial, recreational, scientific, or educational purposes; (C) disease or predation; (D) the inadequacy of existing regulatory mechanisms; and (E) other natural or manmade factors affecting its continued existence. The following is a summary of these factors in relation to Lost River sucker status. For an in-depth treatment of the threats, see the SSA (USFWS 2019).

### **Factor A. The present or threatened destruction, modification, or curtailment of its habitat or range.**

The annual demography monitoring of adult LRS in UKL and the Lost River Units continues the trajectory of decline described by Hewitt et al. (2018), and updated by Krause et al. (2023) with little change in population dynamics (Hewitt, et al. 2018, Hewitt et al. 2021, Krause et al. 2023).

A new negative change in population size and structure occurred in Tule Lake National Wildlife Refuge sumps 1A and 1B. During 2021-2022, the sumps were drained and the remaining LRS were salvaged from the water and relocated to Upper Klamath Lake. In 2023 juvenile SARP fish were stocked in Lower Klamath National Wildlife Refuge, which is adjacent to the Tule Lake sumps, and survival monitoring is ongoing. However due to unreliability of water allocations it is unclear if a population located in this portion of historical habitat can persist. Assessments continue.

Loss and alteration of habitats (including spawning and rearing habitats) were major factors leading to the listing of Lost River sucker and continue to be significant threats to recovery through the lasting effects of the legacy of habitat loss and modification. Suitable habitat has drastically declined due to conversion of wetlands to agricultural use and construction of irrigation and hydroelectric facilities that create barriers, which prevent access to spawning habitat and cause mortality by entraining fish. Loss of ecosystem functions provided by effective wetland habitat continues to threaten the recovery of Lost River sucker through numerous pathways, including effects on water quality, water quantity, access to refugial areas, and food web dynamics.

Prevailing degraded quality of remaining areas exacerbates the effects of pervasive habitat loss. Most water bodies currently occupied by Lost River sucker do not meet water quality standards for nutrients, dissolved oxygen, temperature, and pH set by the States of Oregon and California. These conditions, which manifest primarily in summer, have caused several incidents of widespread adult mortality. The occurrence of mass mortality of fish in Upper Klamath Lake is not new; however, the modern dominance of the cyanobacterium *Aphanizomenon flos-aquae* in the system has led to increased regularity of extreme events. Although conditions are most severe in Upper Klamath Lake and Keno Reservoir, individuals throughout the basin are vulnerable to water-quality-related mortality. There is some indication that cyanobacterial blooms may now be present during the summer in Clear Lake, which, if true, is a distinct change from previous conditions. Assessments are on-going. Degraded water quality conditions may also weaken fish and increase their susceptibility to disease, parasites, and predation. Water quality remains one of the most important factors threatening Lost River sucker existence.

#### **Factor B: Overutilization for commercial, recreational, scientific, or educational purposes**

Overutilization is not currently a threat to the species. Overharvest contributed to declining population levels of Lost River sucker prior to listing, but harvest of any kind has been restricted since 1987. Regulated collections of these species for scientific purposes continue. These collections are not currently a threat to population status. The demographic effects of these collections are regularly evaluated under the ESA Section 7 consultation process (USFWS 2023).

#### **Factor C: Disease or predation**

We identified non-native species as a potential threat at the time of listing through predation or as novel sources or modifiers of diseases and parasites. Several species of birds and fish (native and non-native) prey on Lost River sucker. The ultimate effect to the status of the species from these continues to be investigated and avian predation models are still being refined (Evans et al. 2022). Recent information suggests parasites and disease could be a current or future threat to the Lost River sucker especially at the juvenile life-stage (Markle et al. 2020, J. Lovy

personal comms. 2024).

#### **Factor D: Inadequacy of existing regulatory mechanisms**

Based on the protections provided by existing regulatory mechanisms, we do not consider the inadequacy of regulatory mechanisms to be a threat to the Lost River sucker. Several federal and state regulations directly and indirectly provide protection for Lost River sucker. The primary regulatory authorities are the Endangered Species Acts enacted by the Federal Government and the States of Oregon and California. Lost River sucker are listed as endangered by each of these entities. In general, such authorities prohibit activities that will harm the species' chances of survival and/or recovery, including contemporary protection of habitat.

#### **Factor E. Other natural or manmade factors affecting its continued existence**

##### *Climate Change*

Lost River sucker are adapted to withstand periodic droughts, but given the current reduced state of the species, increases in the intensity or frequency of droughts or a substantial shift in the timing of snowmelt and runoff may negatively affect Lost River sucker. Increasing temperatures could also pose a threat to populations because present-day maximum temperatures already approach stressful ranges for the species. Climate variability results from natural processes; however, climatic models suggest that anthropogenic causes drive much of the recent trends in global climate. It is difficult to predict accurately how such climatic changes will affect the Lost River sucker.

##### *Conservation Efforts*

We, together with many other entities and partners, have accomplished much to reduce threats to Lost River sucker since their listing, much of which is discussed in the revised Recovery Plan and most recent 5-year Status Review (USFWS 2012, 2019). Since the last 5-year Status Review, the KFNFH continues to be our strongest tool to bolster wild populations. With the construction of the hatchery and successful development of captive broodstock, we expect to reach our goal of rearing and repatriating up to 60,000 suckers (of all three species) to UKL annually. Since the inception of the hatchery in 2015, over 70,000 juvenile suckers (of all three species) have been added to augment the wild population.

#### **SYNTHESIS**

The status of the Lost River sucker, relative to ESA listing, has not markedly changed since the previous 5-year Status Review and the SSA in 2019. Overall resiliency for this species is still low and is declining with time as the adult spawning populations reach the end of their expected lifespan. Redundancy is also critically low. There are only three distinct spawning populations: Upper Klamath Lake-springs, Upper Klamath Lake-river, and Clear Lake Reservoir. Two of these spawning populations (Clear Lake Reservoir and Upper Klamath Lake-springs) have very low numbers and are extremely vulnerable to localized catastrophic events, such as fish kills due to poor water quality or inability to access spawning habitat during low water years. The Clear Lake Reservoir population is isolated from the others and adults are often cut off from accessing spawning tributaries due to high evapotranspiration and low water levels. Lost River sucker that spawn at the eastern shoreline springs of Upper Klamath Lake are unique as the only known spawning congregation outside of a river environment. This ecological redundancy is an

important aspect of diversity because it could provide resilience to future localized disturbances. However, juveniles produced from both spawning populations are subject to similar conditions in Upper Klamath Lake, and both experience recruitment failure due to high juvenile mortality.

As a species, Lost River sucker appear to be relatively genetically distinct. Mitochondrial DNA suggests only about 2 percent of Lost River sucker introgress with other species. This is the lowest of all the sucker species within the basin (Dowling et al. 2016 pp. 12 & 13). Nevertheless, the known genetic distinction from shortnose sucker is still relatively low (Hoy and Ostberg 2015 p. 675). Given these conditions, the species has a high degree of the threat of extinction and a low recovery possibility. Therefore, no change in status is warranted at this time.

## RESULTS

### Recommended Classification:

- Downlist to Threatened
- Uplist to Endangered
  - Delist (*Indicate reasons for delisting per 50 CFR 424.11*):
  - Extinction
  - Recovery
  - Original data for classification in error
- No change needed

**New Recovery Priority Number:** No change (4C)

**Brief Rationale:** This is due to a monotypic species with a high degree of the threat of extinction, a low recovery possibility (4), and conflict with economic development (C).

## RECOMMENDATIONS FOR FUTURE ACTIONS

### Establish a Klamath Sucker Recovery Program

This effort was identified as a priority 1 Recovery Action (Action # 8) in the revised Recovery Plan (USFWS 2012). A Recovery Program should be structured to coordinate coherent, and consistent science-based, cooperative actions required to recover species. To do this, a Recovery Program maintains a dedicated staff structure, and coordinates recovery priorities and actions via a clear process of engagement with external partners. Actions from the Recovery Program should focus on (per Recovery Plan Actions 8.1-8.4.2): facilitating information exchange and synthesis; periodically assessing effectiveness of and adjusting recovery actions and priorities; and cause and effect monitoring. As part of these actions, the Recovery Program also identifies, supports, and informs use of recovery tools (e.g. the SARP program, off-channel rearing, etc.).

### Expand and Improve the Rearing Program

The rearing program provides the best short-term avenue to increase the number of Lost River sucker in Upper Klamath Lake. This effort was identified as a priority Recovery Action (Action # 5) in the revised Recovery Plan (USFWS 2012). Actions should focus on improving the survival and growth of individuals while in captivity, maximizing survival and recruitment once reintroduced into Upper Klamath Lake, and increasing the overall numbers of individuals stocked. Development and maintenance of captive brood stock and the refinement of husbandry practices to ensure production of viable larvae is a top priority in planning for potential catastrophic events.

### **Establish at least Two Auxiliary Populations**

The establishment of two auxiliary populations will help the team plan for stochastic events on the landscape and provide a level of redundancy (Recovery Action #5). One location for an auxiliary population could be through the re-establishment of LRS on the landscape at Tule Lake, while a second location is designated and planned for.

### **Phosphorus Reduction**

Poor water quality conditions in Upper Klamath Lake appear to be the most likely driver of persistent recruitment failures in the system. The principal driving factor of the water quality dynamics is the influx of phosphorus into the lake. Specific plans should be developed, and actions implemented to secure long-term reduction of phosphorus levels in the system (Recovery Action #2).

### **Adaptive Management**

The most critical need for this species is to restore natural rates of recruitment to all populations. However, the complexity of biological and physical systems makes it very difficult to determine how to achieve this goal. Adaptive management should be used to implement recovery efforts (including the priorities named above) to ensure the most effective progress towards recovery (Recovery Action #8). Adaptive management includes a cycle of deciding on priority actions, action design, implementation, appropriate monitoring, evaluation, and adjustment of priorities and designs as necessary.

## REFERENCES

- Banet, N.V., and D.A. Hewitt. 2019. Monitoring of endangered Klamath Basin suckers translocated from Lake Ewauna to Upper Klamath Lake, Oregon, 2014–2017: U.S. Geological Survey Open-File Report 2019–1085, 40 p.
- Banet, N.V., D.A. Hewitt, A. Dolan-Caret, and A.C. Harris. 2021. Spatial and temporal distribution of radio-tagged Lost River (*Deltistes luxatus*) and shortnose (*Chasmistes brevirostris*) suckers in Clear Lake Reservoir and associated spawning tributaries, Northern California, 2015–17: U.S. Geological Survey Open-File Report 2021–1061, 37 p.
- Bart, R.J., S.M. Burdick, M.S. Hoy, and C.O. Ostberg. 2020. Juvenile Lost River and shortnose sucker year class formation, survival, and growth in Upper Klamath Lake, Oregon and Clear Lake Reservoir, California 2017: Monitoring Report: U.S. Geological Survey Open-File Report 2020–1025, 36 p.
- Burdick, S.M., C.M. Conway, C.O. Ostberg, R.J. Bart, and D.G. Elliott. 2021. Survival and growth of suckers in mesocosms at three locations within Upper Klamath Lake, Oregon, 2018: U.S. Geological Survey Open-File Report 2021–1036, 18 p.
- Burdick, S.M., D.A. Hewitt, B.A. Martin, L. Schenk, and S.A. Rounds. 2020a. Effects of harmful algal blooms and associated water-quality on endangered Lost River and shortnose suckers. *Harmful Algae* 97:101847.
- Burdick, S.M., D. H. Hereford, C. M. Conway, N.V. Banet, R. Powers, B.A. Martin, and D.G. Elliott. 2020b. Mortality of endangered juvenile Lost River suckers associated with cyanobacteria blooms in mesocosms in Upper Klamath Lake, Oregon. *Transactions of American Fisheries Society* 149:245-265.
- Dowling, T. E., D. F. Markle, G. J. Tranah, E. W. Carson, D. W. Wagman, and B. P. May. 2016. Introgressive Hybridization and the Evolution of Lake-Adapted Catostomid Fishes. *PLoS ONE* 11.
- Evans, A.F., Q. Payton, N. Banet, and B.M. Cramer. 2022. Avian predation on juvenile and adult Lost River and shortnose suckers: an updated multi-predator species evaluation. *American Fisheries Society* 43(6):1561-1574.
- Hereford, D.M., C.M. Conway, S.M. Burdick, D.G. Elliott, T.M. Perry, A. Dolan-Caret, and A.C. Harris. 2019. Assessing causes of mortality for endangered juvenile Lost River suckers (*Deltistes luxatus*) in mesocosms in Upper Klamath Lake, south-central Oregon, 2016: U.S. Geological Survey Open -File Report 2019-1006, 80 p.
- Hewitt, D.A., E.C. Janney, B.S. Hayes, and A.C. Harris. 2018. Status and trends of adult Lost River (*Deltistes luxatus*) and shortnose (*Chasmistes brevirostris*) sucker populations in Upper Klamath Lake, Oregon, 2017: U.S. Geological Survey Open-File Report 2018-1064, 31 p.
- Hewitt, D.A., B.S. Hayes, A.C. Harris, E.C. Janney, C.M. Kelsey, R.W. Perry, and S.M. Burdick. 2021. Dynamics of endangered sucker populations in Clear Lake Reservoir, California: U.S. Geological Survey Open-File Report 2021–1043, 59 p.
- Hoy, M. S., and C. O. Ostberg. 2015. Development of 20 TaqMan assays differentiating the endangered shortnose and Lost River suckers. *Conservation Genetics Resources* 7:673–676.
- Kann, J. and J.D. Walker. 2020. Detecting the effect of water level fluctuations on water quality impacting endangered fish in a shallow, hypereutrophic lake using long-term monitoring

- data. *Hydrobiologia* 847:1851-1872.
- Krause, J.R., E.C. Janney, S.M. Burdick, A.C. Harris, and B.S. Hayes. 2022. Water and endangered fish in the Klamath River Basin: do Upper Klamath Lake surface elevation and water quality affect adult Lost River and shortnose sucker survival? *North American Journal of Fisheries Management* 42:1414-1432.
- Krause, J.R., R.K. Paul-Wilson, and A.C. Harris. 2023. Status and trends of adult Lost River (*Deltistes luxatus*) and shortnose (*Chasmistes brevirostris*) sucker populations in Upper Klamath Lake, Oregon, 2023: U.S. Geological Survey data release.
- Markle, D.F., A. Janik, J.T. Peterson, A. Choudhury, D.C. Simon, V.V. Tkach, M.R. Terwilliger, J.L. Sanders, and M.L. Kent. 2020. Odds ratios and hurdle models: a long-term analysis of parasite infection patterns in endangered young-of-the-year suckers from Upper Klamath Lake, Oregon, USA. *International Journal for Parasitology* 50(4):315-330.
- Martin, B.A., S.M. Burdick, M.K. Purcell, and R.L. Powers. 2021a. Effect of temperature on survival of Lost River suckers with a natural infection of *Ichthyobodo* spp. *North American Journal of Aquaculture* 83:184-191.
- Martin, B.A., S.M. Burdick, S.T. Staiger, and C. Kelsey. 2021b. Water quality, instream habitat, and the distribution of suckers in the upper Lost River watershed of Oregon and California, summer 2018: U.S. Geological Survey Open-File Report 2021-1077, 29 p.
- Martin, B.A., C.M. Kelsey, S.M. Burdick, and R.J. Bart. 2022. Growth, survival, and cohort formation of juvenile Lost River (*Deltistes luxatus*) and shortnose suckers (*Chasmistes brevirostris*) in Upper Klamath Lake, Oregon, and Clear Lake Reservoir, California—2020 monitoring report: U.S. Geological Survey Open-File Report 2022-1099, 27 p.
- Martin, B.A., S.M. Burdick, R.K. Paul-Wilson, and R.J. Bart. 2023. Validating a nonlethal method of aging endangered juvenile Lost River and shortnose suckers. *Journal of Fish and Wildlife Management* 14(1):121-134.
- Smith, M., J. Von Bargen, C. Smith, M. Miller, J. Rasmussen, and D.A. Hewitt. 2020. Characterization of the genetic structure of four sucker species in the Klamath River. USFWS, Abernathy Fish Technology Center, Longview, Washington.
- Smith, M., J. Von Bargen, R. Headley, C. Smith, and J. Spangler. 2023. An improved genetic tool for species identification of Klamath River Basin suckers. USFWS, Abernathy Fish Technology Center, Longview, Washington.
- Walker, J.D., and J. Kann. 2022. Water and nutrient balances of Upper Klamath Lake, water years 1992-2018. Technical Report prepared for the Klamath Tribes Natural Resources Department, Chiloquin, OR. 80 pp + Appendices.
- USFWS. 2012. Revised recovery plan for the Lost River sucker (*Deltistes luxatus*) and shortnose sucker (*Chasmistes brevirostris*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California.
- USFWS. 2019. Species status assessment for the endangered Lost River sucker and shortnose sucker. Klamath Falls, Oregon.
- USFWS. 2023. Biological Opinion on the Effects of the Proposed Interim Klamath Projects Operations Plan, effective October 1, 2023, through October 31, 2024, on the Lost River sucker and shortnose sucker. USFWS Klamath Falls Fish and Wildlife Office, Klamath Falls, Oregon.

**U.S. FISH AND WILDLIFE SERVICE**  
**5-YEAR REVIEW of *species x***

**Current Classification:**

**Recommendation resulting from the 5-Year Review:**

- Downlist to Threatened
- Uplist to Endangered
- Delist (Indicate reasons for delisting per 50 CFR 424.11):
  - Extinction
  - Recovery
  - Original data for classification in error
- No change needed

**Appropriate Listing/Reclassification Priority Number, if applicable:**

**FIELD OFFICE APPROVAL:**

**Lead Field Supervisor, Fish and Wildlife Service**

Approve \_\_\_\_\_ Date \_\_\_\_\_

*The lead Field Office must ensure that other offices within the range of the species have been provided adequate opportunity to review and comment prior to the review's completion. The lead field office should document this coordination in the agency record.*